

CENG 222

Statistical Methods for Computer Engineering

Week 10

Chapter 9

9.4 Hypothesis Testing

Testing Hypotheses

- Hypothesis H_0 and the alternative H_A are two mutually exclusive statements about some unknown parameter θ .
- Testing steps:
 - Collect data
 - Compute a test statistic
 - State if there is sufficient evidence to reject H_0 in favor of H_A
- Examples: 9.22, 9.23, and 9.24

Type I and Type II errors and level of significance

		Result of the test	
		Reject H_0	Accept H_0
H_0 is true	Type I error	correct	
H_0 is false	correct	Type II error	



- We generally want to limit Type I errors (also called false positives) while minimizing Type II errors (also called false negatives)
- Significance level is the probability to observe Type I errors.

Type I and Type II errors and level of significance

- We generally want to limit Type I errors (also called false positives) while minimizing Type II errors (also called false negatives)
- $\alpha = P(\text{reject } H_0 | H_0 \text{ is true})$
- Power of the test is the probability to avoid Type II error
 - Power = sensitivity = recall = True Positive Rate
 - $p(\theta) = P(\text{reject } H_0 | \theta; H_A \text{ is true})$
- See:
 - https://en.wikipedia.org/wiki/Sensitivity_and_specificity


Types of alternatives

- Two sided, one-sided left tail, and one-sided right tail alternatives



Null distribution and acceptance/rejection regions

- The distribution of the test statistic T is called the null distribution.



Z-test

- If the null distribution of the test statistic is Standard Normal, the tests are called Z-tests.
- Z-tests are used when we know population variance.
- T-tests are used for unknown population variance
- Tests can be performed for one sample, two samples (e.g., when comparing two populations)
- Common hypotheses are about population means, proportions, and differences.

Two-tail Z-test

- Data: X_1, \dots, X_n from Normal(μ, σ) with unknown μ and known σ
- Test $H_0 : \mu = \mu_0$ versus $H_A : \mu \neq \mu_0$
 1. Find $\pm z_{\alpha/2}$. Acceptance region is $[-z_{\alpha/2}, z_{\alpha/2}]$
 2. Compute the test statistic
$$Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}}$$
 3. If Z belongs to the acceptance region do not reject H_0
Otherwise, reject H_0

One-sided right tail Z-test

- Data: X_1, \dots, X_n from Normal(μ, σ) with unknown μ and known σ
- Test $H_0 : \mu = \mu_0$ versus $H_A : \mu > \mu_0$
 1. Find z_α . Acceptance region is $(-\infty, z_\alpha]$

2. Compute the test statistic

$$Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}}$$

3. If Z belongs to the acceptance region do not reject H_0
Otherwise, reject H_0

One-sided left tail Z-test

- Data: X_1, \dots, X_n from Normal(μ, σ) with unknown μ and known σ
- Test $H_0 : \mu = \mu_0$ versus $H_A : \mu < \mu_0$
 1. Find z_α . Acceptance region is $[-z_\alpha, +\infty)$

2. Compute the test statistic

$$Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}}$$

3. If Z belongs to the acceptance region do not reject H_0
Otherwise, reject H_0

Unknown variance: T-test

- We use the estimator for variance

$$s^2 = \frac{1}{n - 1} \sum_{i=1}^n (X_i - \bar{X})^2$$

- and use the t -distribution with $n - 1$ degrees of freedom

Z-test summary


Null hypothesis	Parameter, estimator	If H_0 is true:		Test statistic
		$E(\hat{\theta})$	$\text{Var}(\hat{\theta})$	
H_0	$\theta, \hat{\theta}$	$E(\hat{\theta})$	$\text{Var}(\hat{\theta})$	$Z = \frac{\hat{\theta} - \theta_0}{\sqrt{\text{Var}(\hat{\theta})}}$
One-sample Z-tests for means and proportions, based on a sample of size n				
$\mu = \mu_0$	μ, \bar{X}	μ_0	$\frac{\sigma^2}{n}$	$\frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}}$
$p = p_0$	p, \hat{p}	p_0	$\frac{p_0(1-p_0)}{n}$	$\frac{\hat{p} - p_0}{\sqrt{\frac{p_0(1-p_0)}{n}}}$
Two-sample Z-tests comparing means and proportions of two populations, based on independent samples of size n and m				
$\mu_X - \mu_Y = D$	$\mu_X - \mu_Y, \bar{X} - \bar{Y}$	D	$\frac{\sigma_X^2}{n} + \frac{\sigma_Y^2}{m}$	$\frac{\bar{X} - \bar{Y} - D}{\sqrt{\frac{\sigma_X^2}{n} + \frac{\sigma_Y^2}{m}}}$
$p_1 - p_2 = D$	$p_1 - p_2, \hat{p}_1 - \hat{p}_2$	D	$\frac{p_1(1-p_1)}{n} + \frac{p_2(1-p_2)}{m}$	$\frac{\hat{p}_1 - \hat{p}_2 - D}{\sqrt{\frac{\hat{p}_1(1-\hat{p}_1)}{n} + \frac{\hat{p}_2(1-\hat{p}_2)}{m}}}$
$p_1 = p_2$	$p_1 - p_2, \hat{p}_1 - \hat{p}_2$	0	$p(1-p) \left(\frac{1}{n} + \frac{1}{m} \right),$ where $p = p_1 = p_2$	$\frac{\hat{p}_1 - \hat{p}_2}{\sqrt{\hat{p}(1-\hat{p}) \left(\frac{1}{n} + \frac{1}{m} \right)}}$ where $\hat{p} = \frac{n\hat{p}_1 + m\hat{p}_2}{n+m}$

T-test summary

Hypothesis H_0	Conditions	Test statistic t	Degrees of freedom
$\mu = \mu_0$	Sample size n ; unknown σ	$t = \frac{\bar{X} - \mu_0}{s/\sqrt{n}}$	$n - 1$
$\mu_X - \mu_Y = D$	Sample sizes n, m ; unknown but equal standard deviations, $\sigma_X = \sigma_Y$	$t = \frac{\bar{X} - \bar{Y} - D}{s_p \sqrt{\frac{1}{n} + \frac{1}{m}}}$	$n + m - 2$
$\mu_X - \mu_Y = D$	Sample sizes n, m ; unknown, unequal standard deviations, $\sigma_X \neq \sigma_Y$	$t = \frac{\bar{X} - \bar{Y} - D}{\sqrt{\frac{s_X^2}{n} + \frac{s_Y^2}{m}}}$	Satterthwaite approximation, formula (9.12)

Confidence intervals versus two-sided level- α tests

- A level- α two sided test is the same as testing whether a given test statistic is in the $(1 - \alpha)100\%$ confidence interval $[a,b]$



- Examples: 9.31, 9.35 (one-sided test)

P-value

- Instead of a fixed significance level α , we can compute the boundary level for acceptance/rejection for the computed test statistic
 - This computed value is called the p-value
 - It is the probability that another sample results in a more extreme test static.

